


FORTNIGHTLY REPORT OF SEASONAL CONDITION

Integrated Seasonal Condition Monitoring System





Team

Telangana State Remote Sensing Applications Centre (TRAC)

Sri K. Ramakrishna Rao, IAS

Principal Secretary (Finance and Planning)
Government of Telangana
Director General (TRAC)

Dr. G. Sreenivasa Reddy

Addl. Director General

Dr. J. Ramana murthy

Technical Adviser

Sri. G. Mehar Baba

Technical Adviser

Dr. M. Kavitha

Assistant Scientific Officer

Sri. A. Kamalakar Reddy

Assistant Scientific Officer

Sri. N. Narender

Scientific Assistant

Acknowledgement


We take this opportunity to express our sincere thanks to Directorate of Economics and Statistics, Telangana State Developing Planning Society, and India Meteorological Department for providing rainfall data. We also express our gratitude to Department of Agriculture and Irrigation Department for sharing progress of crop sowings and reservoir water levels data respectively for integrated seasonal condition monitoring of the state.

HIGHLIGHTS


INTEGRATED SEASONAL CONDITION MONITORING SYSTEM (ISMS) - TELANGANA

Cumulative Report up to 15th June, 2019


- Seasonal condition is categorised as “Normal” in **171** Mandals as on date 15th June 2019
- Seasonal condition is categorised as “Watch” in **355** Mandals as on date 15th June 2019
- Seasonal condition is categorised as “Alert” in **63** Mandals as on date 15th June 2019



Seasonal Condition up to 1st fortnight of June, 2019



Rainfall from 1st June to 15th June 2019



Seasonal condition of Telangana up to 1st fortnight of June 2019

Rainfall 01st June to 15th June, 2019

- **151** Mandals out of 589 (**26%**) of state received **Deficient** rainfall. **25** Mandals (**4%**) of the state received **Excess** rainfall. **288** Mandals (**49%**) of the state received **Large Deficient** rainfall. **25** Mandals (**4%**) of the state received **Large Excess** rainfall.
- **60** Mandals (**10%**) have received **Normal** rainfall respectively.

CONTENTS

S. No.	Description	Page No
1	Background and Rationale	1
2	Data used, Indicators and Methodology	3
3	Present status up to 1 st fortnight of June 2019	7
3.1	Rainfall data	7
3.2	Reservoir Water Levels	10
3.3	Vegetation Index	11
3.4	Surface Wetness Indicators	14
3.5	Drought situation of Mandals	17
3.6	District Wise NDVI / NDWI / VCI Status	22
4	References	23

List of Tables

Table No.	Description	Page No
1	Classification of agricultural situation	3
2	Data source and indicators	3
3	Rainfall status as on 15 th June 2019	8
4	Reservoir water levels	10
5	Mandals under Watch and Alert category based on ISMS criteria	18
6	District wise NDVI / NDWI / VCI Status	22

List of Figures

Figure No.	Description	Page. No
1	Location of Automatic Weather Stations	2
2	Flow chart of drought assessment methodology	6
3	Deviation of rainfall in percent w.r.t. normal from June 01 st to June 15 th , 2019	9
4	NDVI - MODIS: First Fortnight of June 2019	11
5	NDVI - MODIS, Fortnightly agricultural situation from June 2019, 2018 and 2017	12
6	NDVI deviation (MODIS - 250m), First Fortnight of June 2019 w.r.t. 2013	13
7	NDWI - MODIS: First Fortnight of June 2019	14
8	NDWI - MODIS, Fortnightly agricultural situation from June 2019, 2018 and 2017	15
9	NDWI deviation (MODIS - 250m), First Fortnight of June 2019 w.r.t. 2013	16
10	Mandal wise drought assessment based on ISMS criterion	17

1. Background and Rationale

Drought is a complex natural hazard. It is defined as any deficiency of water to satisfy the normal need to agriculture, livestock, industry, or human population. Drought assessment and monitoring is essential for the agricultural sector to take appropriate mitigation measures. Drought indices derived from satellite data play a major role in assessing the health and condition of the crops/vegetation.

National Agricultural Drought Assessment and Monitoring System (NADAMS) project of National Remote Sensing Centre (NRSC), Indian Space Research Organization (ISRO) established a remote sensing based drought assessment protocol utilizing the Normalized Difference Vegetation Index (NDVI) and Normalized Difference Water Index (NDWI). The Government of India has established Mahalanobis National Crop Forecast Centre (MNCFC) under Department of Agriculture and Cooperation, New Delhi for carrying out drought assessment at national level.

The Department of Agriculture and Cooperation, Government of India published a drought manual in 2016 which suggested parameters like rainfall deficiency, area under sowing, NDVI, NDWI, Moisture Adequacy Index (MAI) and other indicators to declare drought. State Government monitor drought by obtaining information from various sources on key variables of drought which include rainfall, reservoir / lake water levels, surface water / groundwater, soil moisture and sowing / crop conditions etc. The key variables for monitoring drought in Telangana are:

- Meteorological Data - Rainfall and other parameters like Temperature, Wind speed and Relative Humidity (AWS data)
- Weather forecast - Short, medium, extended range
- Soil Moisture (Moisture Adequate Index)
- Sown Area / Crop Health / Stress
- Satellite based Vegetation Index (NDVI/NDWI)
- Stream Flow - Discharge
- Groundwater Levels
- Reservoir and Lake Storage / Level
- Impacts - distress sale and migration of cattle, human migration, fodder availability, drinking water, animal health, employment opportunities in agriculture sector

An extensive weather observation network of 1044 Automatic Weather Stations (AWS) is established in Telangana. Telangana State Development Planning Society (TSDPS) monitors the data and maintains the networks. Figure 1 showing the location of AWS stations in Telangana.

Government of Telangana
Automatic Weather Stations (AWS)




Figure 1: Location of automatic weather stations

Telangana State Remote Sensing Applications Centre (TRAC) has established a protocol ***Integrated Seasonal Condition Monitoring System (ISMS)***. The objectives of the ISMS are

- *Concurrent monitoring of seasonal conditions using remote sensing, extensive weather network data and continuous ground truth.*
- *Develop an early warning (monitoring and forecasting) of drought using suite of indicators, which will help to increase drought preparedness, and identify and implement appropriate Disaster Risk Reduction (DRR) measures.*
- *Early warning to the Districts/Mandals.*

ISMS uses the rainfall data provided by Directorate of Economics & Statistics (DE&S), weekly progress of crop area sowings, groundwater level and its fluctuation, command and non-command area, water releases data, reservoir levels in addition to the Normalized Difference Vegetation Index (NDVI) and Normalized Difference Water Index (NDWI) based methodology of MNCFC. This output is verified through ground truth, additionally in context of the state specific drought declaration criteria. The agricultural situation is classified in three to four categories as per the NRSC i.e. Normal, Watch, Alert for June to August and Normal, Mild, Moderate and Severe for September to October. The details of the classification of agricultural situation are given in Table 1.

Table. 1. Classification of agricultural situation

Duration	Condition	Description
July - August	Normal	<ul style="list-style-type: none"> • Agricultural situation is normal
	Watch	<ul style="list-style-type: none"> • Progress of agricultural situation is slow • Ample scope for recovery • No external intervention needed
	Alert	<ul style="list-style-type: none"> • Very slow progress of agricultural situation • Need for intervention. • Develop and implement contingency plans to minimise loss
September - October	Mild drought	<ul style="list-style-type: none"> • Crops have suffered stress slightly
	Moderate drought	<ul style="list-style-type: none"> • Considerable loss in production. • Take measures to alleviate suffering
	Severe	<ul style="list-style-type: none"> • High risk significant reduction in crop yield • Management measures to provide relief

2. Data used, Indicators and Methodology

2.1. Data used

Details of data used under project are discussed in Table 2.

Table. 2. Data source and indicators

Data source	Product	Indicators
MODIS (250/500m)	Surface reflectance	NDVI & NDWI
AWiFS	Surface reflectance	NDVI & NDWI
AWS/ DES	<ul style="list-style-type: none"> • Daily rainfall • Crop sown area • Crop cutting experiments 	<ul style="list-style-type: none"> • Rainfall deviation • Dry spells • Crop yield
Agriculture Department, GoTS	Weekly sowing progress	District wise sown areas deviation from normal
Irrigation Department, GoTS	Reservoir levels/ Water release data	Command area Mandals under canal irrigation

2.2. Indicators and Index

2.2.1. Rainfall data

In Telangana, South-West Monsoon is crucial for agriculture sector. ISMS uses integrated (AWS+DES+IMD) Mandal wise rainfall data provided by Directorate of Economics & Statistics (DES). This data is used for computation of meteorological drought situation and to derive the mandal wise spatial distribution of rainfall in the state.

2.2.2. Reservoir water levels and water release - major and medium project

A scheme having Culturable Command Area (CCA) up to 2,000 hectares individually is classified as minor irrigation scheme. A scheme having CCA more than 2,000 hectares and up to 10,000 hectares individually is a medium irrigation scheme. A scheme having CCA more than 10,000 hectares is major irrigation scheme. In Telangana, water is released during Kharif season to major and medium command areas.

2.2.3. Crop sowing progress

Weekly crop sowing progress reports are taken from 'Season and Crop Coverage Report-Kharif 2019' of Commissioner of Agriculture, Telangana. The report includes current status of Weather condition, Water level, Crop sowing and Agricultural Operations.

2.2.4. Vegetation index

The crop/vegetation reflects high energy in the near infrared band due its canopy geometry and health of the standing crops/vegetation and absorbs high in the red band due to its biomass and photosynthesis. Uses of these contrast characteristics of vegetation in near infrared and red bands indicate both the health and condition of the crops/vegetation. Normalised Difference Vegetation Index (NDVI) is widely used for operational drought assessment because of its simplicity in calculation, easy to interpret and its ability to partially compensate for the effects of atmosphere, illumination geometry etc., (Malingreau 1986, Tucker and Chowdhary 1987, Kogan 1995). NDVI is derived by the difference of these measurements and divided by their sum.

$$NDVI = \frac{(NIR - Red)}{(NIR + Red)}$$

The vegetation index is generated from each of the available satellite data irrespective of the cloud cover present. To minimize the cloud, monthly time composite vegetation index is generated.

2.2.5. Surface wetness indicator

Shortwave Infrared (SWIR) band is sensitive to moisture available in soil as well as in crop canopy. In the beginning of the cropping season, soil background is dominant hence SWIR is sensitive to soil moisture in the top 1-2 cm. As the crop progresses, SWIR becomes sensitive to leaf moisture content. SWIR band provides only surface wetness information. When the crop is grown-up, SWIR response is only from canopy and not from the underlying soil. NDWI using SWIR can complement NDVI for drought assessment particularly in the beginning of the cropping season. NDWI is derived as under;

$$NDWI = \frac{(NIR - SWIR)}{(NIR + SWIR)}$$

Higher values of NDWI signify more surface wetness. The wetness index is generated from each of the available satellite data irrespective of the cloud cover present. To minimize the cloud, monthly time composite wetness index is generated.

2.2.6. Vegetation condition index

Kogan (1995) developed Vegetation Condition Index (VCI) using the range of NDVI as under,

$$VCI = \frac{(NDVI - NDVI \min)}{(NDVI \max - NDVI \min)} * 100$$

The current drought assessment expressed as percentage of deviation of NDVI and NDWI based on 10 year NDVI and NDWI index values. The minimum and maximum value of NDVI and NDWI, the VCI discriminated between the weather components.

2.3 Methodology




Figure 2: Flow chart of drought assessment methodology

The methodology to assess and monitor the agricultural conditions and situation in the state at district and Mandal level uses IRS Resourcesat-2 AWiFS data. Indian Remote Sensing satellite (IRS) Resourcesat-2 having Advanced Wide Imaging Field Sensor (AWiFS) payload collects data in two spectral bands 0.62-0.68 μm (red) and 0.77-0.86 μm (near infrared) with spatial resolution of 56 m and ground swath of 740 km with a revisit period of 5 days. Along with this MODIS 250/500 m satellite data provide spectra, radiometric and spatial resolutions products for better monitoring of the agriculture. The combination of AWiFS and MODIS is useful to increase the frequency of images.

The different activities carried out through ISMS commence with acquisition of MODIS (250 m) and AWiFS (56 m) satellite data. The satellite data being processed and NDVI and NDWI indices are developed. Based on these indices deviation with respect to normal year (2013) is calculated and Mandal wise statistics are derived. The agricultural situation is assessed incorporating rainfall deviation, command and non command areas, dry spell, drought prone border line areas, crop sown area progress and ground truth along with satellite derived indices. The flow chart of methodology is shown in Figure 2.

3. Present status up to 1st fortnight of June 2019

3.1. Rainfall data

The status of rainfall as on 15th June 2019 is shown in Table.3.

- **25** Mandal (**4%**) of the state received **Large Excess** (+60% and above) rainfall.
- **25** Mandals (**4%**) of the state received **Excess** (+20% to +59%) rainfall.
- **60** Mandals (**10%**) have received **Normal** (+19% to -19%) rainfall.
- **151** Mandals out of 589 (**26%**) of state received **Deficient** (-20% to -59%) rainfall.
- **288** Mandals (**49%**) of the state received **Large Deficient** (-60% to -99%) rainfall.

Table. 3. Rainfall status as on 15th June 2019

S. No	District Name	Large Excess	Excess	No Rain	Normal	Deficient	Large Deficient	Total
1	Adilabad		1			1	16	18
2	Bhadradri Kothagudem		1		1	4	17	23
3	Hyderabad		2		4	9	1	16
4	Jagtial		2		2	12	2	18
5	Jangoan					5	7	12
6	Jayashankar Bhupalpally			3	1	1	6	11
7	Jogulamba Gadwal	4	1	1	2	2	2	12
8	Kamareddy			1	2	4	15	22
9	Karimnagar		2	1		4	9	16
10	Khammam			9		2	10	21
11	Kumurambheem Asifabad					4	11	15
12	Mahabubabad			5		4	7	16
13	Mahabubnagar	5	3		5	2		15
14	Mancherial			2		5	11	18
15	Medak				2	8	10	20
16	Medchal Malkajgiri				3	8	4	15
17	Mulugu			3			6	9
18	Nagarkurnool	6	2		6	3	3	20
19	Nalgonda		2	3	4	4	18	31
20	Narayanpet		2	1	5	2	1	11
21	Nirmal			1		1	17	19
22	Nizamabad			1		1	27	29
23	Peddapalli		1	1	1	2	9	14
24	Rajanna Sircilla					2	11	13
25	Rangareddy	4	3		6	11	3	27
26	Sangareddy		1		4	11	10	26
27	Siddipet		1		2	8	12	23
28	Suryapet			1	1	8	13	23
29	Vikarabad	3	1		5	6	3	18
30	Wanaparthy	3			3	6	2	14
31	Warangal Rural			5		3	8	16
32	Warangal Urban			2			9	11
33	Yadadri Bhongir				1	8	8	17
		25	25	40	60	151	288	589

SOURCE: DE&S




Figure 3: Deviation of rainfall in percent w.r.t. normal from June 01st to June 15th, 2019

3.2. Reservoir water levels

All the major reservoirs are holding 211 TMC as on 15-06-2019, and as on date last year the level had stood at 353 TMC. The details of water levels of all major reservoirs as on 15-06-2019 are furnished hereunder in Table.4.

Table.4. Reservoir Water Levels

PARTICULARS OF MAJOR RESERVOIRS AS ON 15/June /2019										
SI No	Reservoir Name	Time	FRL (feet)	Gross Capacity (TMC)	THIS YEAR As on 15 / June / 2019				LAST YEAR As on 15 / June / 2018	
					Level	Gross Storage	Inflow	Outflow	Level	Gross Storage
					(in feet)	(TMC)	(Cusecs)	(Cusecs)	(in feet)	(TMC)
Krishna Basin										
1	Almatti	00:00	1705	129.721	0	0	0	0	1667.06	22.41
2	Jurala	09:47	1045	9.657	1027.26	2.15	0	174	1036.91	5.26
3	Nagarjunasagar	07:33	590	312.045	508.3	128.803	1399	1399	511.6	134.403
4	Narayanapur	00:00	1615	37.646	0	0	0	0	1604.43	24.45
5	Srisailam	09:48	885	215.807	806.3	32.24	0	252	800	28.98
6	Tungabhadra	09:48	1633	100.86	1574.56	2.23	0	231	1592.57	11.91
7	Ujjaini	09:47	1630	117.24	1592.55	33.36	0	217	1605.99	53.76
Godavari Basin										
8	Jaikwad	09:50	1522	102.732	1490.29	20	0	648	1503.67	43.69
9	Kaddam	09:52	700	7.6	671.375	2.344	0	70	683.53	4.051
10	Lower Manair Dam	09:51	920	24.074	882.75	3.75	0	220	882	3.56
11	Nizam sagar	09:50	1405	17.803	1364.84	0.03	0	10	1385.14	2.41
12	Singur	09:50	1717.93	29.91	1669.34	0.36	0	90	1697.42	7.92
13	Sri Ram Sagar	09:51	1091	90.313	1049	5.66	0	300	1056.5	10.392

Source: Irrigation Department, Hyderabad

3.3. Vegetation index

The Normalized Difference of Vegetation Index (NDVI) for 1st fortnight of June 2019 is shown in the figures and also compared with 2018 and 2017. The year 2013 is treated as a normal year. Mandal wise NDVI, monthly agricultural situation for the year 2019, 2018 and 2017, deviation of NDVI w.r.t. 2013 are shown in the Figures 4, 5, and 6 respectively. As per NDVI deviation w.r.t normal moderate stress is observed in few Districts.




Figure 4: NDVI - MODIS: First Fortnight of June 2019

Government of Telangana
Normalized Difference Vegetation Index
MODIS (250m) Mandal Wise




Figure 5: NDVI - MODIS, Fortnightly agricultural situation from June 2019, 2018 and 2017




Figure 6: NDVI deviation (MODIS - 250m), First Fortnight of June 2019 w.r.t. 2013

3.4. Surface wetness indicator

The map indicates status of moisture availability in soil as well as in crop canopy for 1st fortnight of June 2019. The year 2013 is treated as a normal year. Mandal wise Normalized Difference Water Index (NDWI) situation the year 2019, 2018 & 2017, Fortnightly agricultural situation deviation of NDWI w.r.t. 2013 are shown in the Figures 7, 8, and 9 respectively. As per NDWI deviation w.r.t normal moderate stress is observed in few Districts.




Figure 7: NDWI - MODIS: First Fortnight of June 2019

Government of Telangana
Normalized Difference Water Index
MODIS (250m) Mandal Wise




Figure 8: NDWI - MODIS, Fortnightly agricultural situation from June 2019, 2018 and 2017




Figure 9: NDWI deviation (MODIS - 250m), First Fortnight of June 2019 w.r.t. 2013

3.5. Drought situation of Mandals

3.5.1 Composite criteria

The drought situation in the state is assessed using different indicators viz., NDVI, NDWI and rainfall deviation of mandals. Compositing all indicators, mandals were categorised into Normal, Watch and Alert. Mandal-wise analysis for the 1st fortnight of June 2019 indicated “Normal” agricultural situation in **171** Mandals. The agricultural situation is categorized as “**Watch**” in **355** and “**Alert**” in **63** Mandals. The Mandals under Watch and Alert categories are given in the Table.5 and their spatial distribution is shown in Figure 10.




Figure 10: Mandal wise drought assessment based on ISMS criterion

Table.5. Mandals under Watch and Alert category based on ISMS criteria

District	Watch (355)	Alert (63)
Adilabad	Total: 08 Adilabad Urban, Bazarhathnoor, Bheempoor, Boath, Ichoda, Mavala, Sirikonda, Utnur.	Total: 09 Adilabad Rural, Bela, Gadiguda, Gudihathnur, Inderavelly, Jainad, Narnoor, Neradigonda, Talamadugu.
Bhadradri Kothagudem	Total: 18 Allapalli, Annapureddipalle, Aswapuram, Burgampadu, Chandrugonda, Cherla, Chunchupally, Dummugudem, Gundala, Julurupad, Karakagudem, Laxmidevipally, Manuguru, Mulakalapally, Palvancha, Pinapaka, Sujathanagar, Yellandu.	Total: 02 Kothagudem, Tekulapalle.
Jagtial	Total: 07 Buggaram, Ibrahimpatnam, Korutla, Mallapur, Medipalle, Raikal, Sarangapur.	
Jangaon	Total: 09 Chilpur, Ghanpur (Stn), Jangoan, Kodakandla, Lingalaghanpur, Narmetta, Ragunathpally, Tharigoppula, Zaffergadh.	
Jogulamba	Total: 04 Alampur, Dharoor, Kaloor Thimmandoddi, Rajoli.	
Jayashankar Bhupalpally	Total: 10 Bhupalpalle, Chityal, Mangapet, Mogullapalle, Mulug, Palmela, Regonda, Tadvai, Tekumatla, Venkatapur.	
Kamareddy	Total:09 Bhiknoor, Bibipet, Bichkunda, Birkoor, Domakonda, Kamareddy, Lingampet, Machareddy, Tadvai.	Total: 11 Banswada, Gandhari, Jukkal, Madnur, Nasurullabad, Nizamsagar, Pedda Kodapgal, Pitlam, Rajampet, Sada-shivnagar, Yellareddy.
Karimnagar	Total: 10 Ellandakunta, Huzurabad, Jammikunta, Karimnagar, Karimnagar Rural, Kothapally, Manakondur, Thimmapur, V Saidapur, Veenavanka.	
Khammam	Total: 20 Bonakal, Chintakani, Enkoor, Kalluru, Kamepally, Khammam_Urban, Konijerla, Kusumanchi, Madhira, Mudigonda, Nelakondapally, Penuballi, Raghunadhapalem, Sathupally, Singareni, Thallada, Tirumalayapalem, Vemsoor, Wyra, Yerrupalem.	

Komaram Bheem	Total: 13 Asifabad, Bejjur, Dahegaon, Jainoor, Kagaznagar, Kerameri, Kouthala, Lingapur, Penchikalpet, Sirpur(T), Sirpur(U), Tiryani, Wankidi.	
Mahabubabad	Total: 13 Bayyaram, Chinnagudur, Dornakal, Gangaram, Garla, Gudur, Kesamudram, Kothaguda, Mahabubabad, Maripeda, Narsimhulapet, Nellikudur, Thorur.	Total: 01 Kuravi.
Mahabubnagar	Total: 02 Gandeed, Nawabpet.	
Mancherial	Total: 14 Bellampalle, Bheemaram, Bheemini, Chennur, Dandepalle, Jaipur, Jannaram, Kannepalli, Kotapalle, Mancherial, Mandamarri, Naspur, Nennel, Vemanpalle.	
Medak	Total: 13 Alladurg, Chilipched, Havelighanpur, Kowdipally, Medak, Narsapur, Narsingi, Nizampet, Papannapet, Ramayampet, Shankarampet(A), Shankarampet(R), Yeldurthy.	Total: 01 Regode.
Medchal Malkajgiri	Total: 09 Alwal, Bachupally, Dundigal Gandimaisamma, Ghatkesar, Keesara, Malkajgiri, Medchal, Muduchinthalphally, Shamirpet.	
Mulugu	Total: 09 Eturnagaram, Govindaraopet, Kannaigudem, Mangapet, Mulug, Tadvai (Sammakka Sarakka), Venkatapur, Venkatapuram, Wazeed.	
Nagarkurnool	Total: 04 Bijinapalle, Kollapur, Lingal, Pentlavelli.	
Nalgonda	Total: 24 Adavidevulapally, Anumula_Haliya, Chandur, Chinthapally, Damaracherla, Gundlapally, Gurrampode, Kanagal, Kethepally, Kondamallapally, Madugulapally, Marriguda, Miryalaguda, Nakrekal, Nalgonda, Nampally, Neredugommu, Nidamanoor, Peda Adisharla Palli, Peddavoora, Tipparthy, TirumalagiriSagar, Tripuraram, Vemulapally.	Total: 01 Chityal.
Narayanpet	Total: 02 Krishna, Marikal.	

Nirmal	Total: 14 Dasturabad, Dilawarpur, Kaddampeddur, Kuntala, Laxmanchanda, Lokeshwaram, Mamda, Narsapur(G), Nirmal, Nirmal Rural, Pembi, Sarangapur, Soan, Tanoor.	Total: 05 Basar, Bhainsa, Khanapur, Kubeer, Mudhole.
Nizamabad	Total: 19 Armoor, Balkonda, Bheemgal, Dharpally, Dichpally, Indalwai, Jakranpally, Kammarpally, Makloor, Morthad, Mugpal, Mupkal, Nandipet, Nizamabad North, Nizamabad Rural, Nizamabad South, Sirikonda, Vailpur, Yergatla.	Total: 09 Bodhan, Chandur, Kotagiri, Mosra, Navipet, Renjal, Rudrur, Varni, Yedapally.
Peddapalli	Total: 12 Anthergaon, Eligaid, Julapalli, Kamanpur, Manthani, Mutharam_Manthani, Odela, Peddapalli, Ramagiri, Ramagundam, Srirampur, Sultanabad.	
Rajanna Sirsilla	Total: 10 Chendurthi, Gambhiraopet, Konaraopet, Mustabad, Rudrangi, Sircilla, Thangallapalli, Veernapalli, Vemulawada Rural, Yellareddypet.	Total: 01 Vemulawada.
Rangareddy	Total: 05 Chevella, Gandipet, Moinabad, Shamshabad, Shankarpalle.	Total: 03 Chowdergudem, Kondurg, Rajendranagar.
Sangareddy	Total: 10 Gummadidala, Kalher, Kandi, Kondapur, Nagalgidda, Patancheruvu, Pulkal, Raikode, Sangareddy, Vatpally.	Total: 10 Jharasangam, Kangti, Kohir, Mogudampally, Muni-pally, Narayankhed, Nyalkal, Sadasivpet, Sirgapor, Zahirabad.
Siddipet	Total: 14 Akkannapet, Bejjanki, Cherial, Chinnakodur, Dubbak, Jagdevpur, Maddur, Markook, Mirdoddi, Narayanraopet, Raipole, Siddipet Rural, Siddipet Urban, Thoguta.	
Suryapet	Total: 17 Ananthagiri, Chilkur, Chinthala palem, Garidepalli, HuzurNagar, Kodad, Maddirala, Mattampalli, Mellachervu, Mothey, Munagala, Nadigudem, Nereducherla, Palakeedu, Penpahad, Suryapet, Thungathurthy.	
Vikarabad	Total: 08 Bommaraspeta, Dharoor, Doma, Kotepally, Pargi, Peddemul, Pudur, Vikarabad.	Total: 04 Bantwaram, Kulkacharla, Marpalle, Mominpet.

Wanaparthy	Total: 05 Atmakur, Chinnambavi, Pebbair, Srirangapur, Weepangandla.	
Warangal Rural	Total: 15 Atmakur, Chennaraopet, Duggondi, Geesugonda, Khanapur, Nadikuda, Nallabelli, Narsampet, Nekkonda, Parkal, Parvathagiri, Raiparthy, Sangem, Shayampet, Wardhannapet.	Total: 01 Damera.
Warangal Urban	Total: 11 Bheemadevarapalli, Dharmasagar, Elkathurthi, Hanamkonda, Hasanparthy, Inavolu, Kamalapur, Khaazipet, Khila Warangal, Velair, Warangal.	
Yadadri Bhongir	Total: 12 Alair, Athmakur (M), B.Pochampally, Bhongiri, Bibinagar, Bommalaramaram, Choutuppal, Mothkur, Rajapeta, Ramannapet, Valigonda, Yadagirigutta.	

3.6. District Wise NDVI / NDWI / VCI Status

NDVI/NDWI/VCI status as on 15/06/2019, Telangana								
S.No	District	NDVI Value	Average NDVI	NDWI Value	Average NDWI	VCI (NDVI)	VCI (NDWI)	VCI Condition
1	Adilabad	0.219	0.235	0.017	0.026	26.26	16.28	Moderate
2	Bhadradri-Kothagudem	0.317	0.337	0.046	0.107	51.86	16.76	Mild
3	Hyderabad	0.221	0.250	0.099	0.103	40.63	46.77	Mild
4	Jagtial	0.256	0.266	0.064	0.076	44.66	37.18	Mild
5	Jangaon	0.262	0.230	0.029	0.022	65.22	24.89	Mild
6	Jayashankar-Bhupalpally	0.254	0.240	0.038	0.031	58.25	52.60	Mild
7	Jogulamba-Gadwal	0.245	0.229	0.044	0.020	51.59	11.72	Mild
8	Kamareddy	0.237	0.264	0.020	0.061	36.72	5.28	Moderate
9	Karimnagar	0.236	0.212	0.058	0.059	62.98	45.67	Normal
10	Khammam	0.339	0.343	0.060	0.114	61.61	26.68	Mild
11	Komaram Bheem-Asifabad	0.232	0.217	0.027	0.036	55.92	17.69	Mild
12	Mahabubabad	0.277	0.292	0.025	0.053	53.95	10.55	Mild
13	Mahabubnagar	0.246	0.251	0.038	0.033	48.77	13.26	Mild
14	Mancherial	0.258	0.220	0.046	0.036	67.11	52.27	Normal
15	Medak	0.236	0.258	0.039	0.050	42.97	14.49	Mild
16	Medchal-Malkajgiri	0.274	0.272	0.050	0.081	54.77	17.56	Mild
17	Mulug	0.273	0.254	0.029	0.057	63.45	24.60	Mild
18	Nagarkurnool	0.233	0.213	0.049	0.017	53.95	13.35	Mild
19	Nalgonda	0.239	0.234	0.042	0.043	59.70	18.12	Mild
20	Narayanpet	0.226	0.222	0.058	0.019	55.72	6.91	Mild
21	Nirmal	0.240	0.251	0.036	0.044	40.09	30.45	Mild
22	Nizamabad	0.226	0.234	0.021	0.047	50.07	19.83	Mild
23	Peddapalli	0.235	0.243	0.073	0.072	49.65	47.82	Mild
24	Rajanna-Siricilla	0.237	0.217	0.019	0.018	61.78	26.86	Mild
25	Rangareddy	0.256	0.267	0.039	0.059	45.17	16.75	Mild
26	Sangareddy	0.245	0.280	0.043	0.095	27.12	9.57	Moderate
27	Siddipet	0.247	0.223	0.033	0.021	64.88	14.46	Moderate
28	Suryapet	0.253	0.250	0.042	0.056	62.80	15.91	Moderate
29	Vikarabad	0.236	0.280	0.027	0.082	25.09	4.13	Moderate
30	Wanaparthy	0.247	0.218	0.015	0.044	56.34	7.96	Mild
31	Warangal Rural	0.265	0.286	0.020	0.059	44.01	23.17	Mild
32	Warangal Urban	0.263	0.264	0.039	0.069	54.53	30.27	Mild
33	Yadadri-Bhongir	0.252	0.227	0.029	0.038	64.77	11.07	Moderate

Table.6 District wise NDVI / NDWI / VCI Status

*Normalized Difference Vegetative Index (NDVI) Value - Current year NDVI

*Normalized Difference Wetness Index (NDWI) Value - Current year NDWI

*Average NDVI - Average of previous 16 years NDVI

*Average NDWI - Average of previous 16 years NDWI

*VCI (NDVI) - Vegetation Condition Index based on NDVI

*VCI (NDWI) - Vegetation Condition Index based on NDWI

*NDVI/NDWI Condition - VCI>=60 (Normal), VCI>=40 (Mild), VCI>=20 (Moderate), VCI<20 (Severe)

4. References

Department of Agriculture and Cooperation, 2009, Manual for Drought Management, Ministry of Agriculture, Govt. of India, New Delhi.

<http://drought.unl.edu/portals/0/docs/international/GovtIndiaDroughtManual.pdf>

Department of Agriculture, 2017, Season and Crop Coverage Report, Kharif - 2017, Govt. of Telangana

Kogan FN, 1995, Droughts of late 1980s in the USA as derived from NOAA polar orbiting satellite data, Bulletin of American Meteorological Society, 76: 655-668

Malingreau JP, 1986, Global vegetation dynamics: Satellite observations over Asia, International Journal of Remote Sensing, 7: 1121-1146.

Tucker CJ and Chowdhary BJ, 1987, Satellite remote sensing of drought conditions, Remote Sensing of Environment, 23: 243-251

